

Fractional Pearson Diffusions

N. Leonenko¹

¹ Cardiff School of Mathematics, Cardiff University,
Senghennydd Road, Cardiff CF24 4YH, United Kingdom

Abstract

Pearson diffusions have stationary distributions of Pearson type. They includes Ornstein-Uhlenbeck, Cox-Ingersoll-Ross, and several others well-known processes. Their stationary distributions solve the Pearson equation, developed by Pearson in 1914 to unify some important classes of distributions (e.g., normal, gamma, beta). Their eigenfunction expansions involve the traditional classes of orthogonal polynomials (e.g., Hermite, Laguerre, Jacobi). We develop fractional Pearson diffusions ([1],[2]), constructing by a non-Markovian inverse stable time change. Their transition densities are shown to solve a time-fractional analogue to the diffusion equation with polynomial coefficients. Because this process is not Markovian, the stochastic solution provides additional information about the movement of particles that diffuse under this model. This is joint work with M.M. Meerschaert and A. Sikorskii (Michigan State University, USA).

Anomalous diffusions have proven useful in applications to physics, geophysics, chemistry, and finance.

REFERENCES

[1] Leonenko, N.N., Meerschaert, M.M and Sikorskii, A. (2013) Fractional Pearson diffusion, *Journal of Mathematical Analysis and Applications*, vol. 403, 532-546

[2] Leonenko, N.N., Meerschaert, M.M and Sikorskii, A. (2013) Correlation Structure of Fractional Pearson diffusion, *Computers and Mathematics with Applications*, 66, 737-745