

A REPRESENTATION FOR THE DERIVATIVE WITH RESPECT TO THE INITIAL DATA OF THE SOLUTION OF AN SDE WITH NON-REGULAR DRIFT

OLGA ARYASOVA

We consider a multidimensional SDE with an identity diffusion matrix and a drift vector being a vector function of bounded variation.

According to [8] there exists a unique strong solution to such an equation.

It is well known that if the drift vector is continuously differentiable and its derivative is bounded then the SDE generates a flow of diffeomorphisms. It turns out that this condition can be essentially reduced [5], and a flow of diffeomorphisms exists in the case of possible unbounded Hölder continuous drift vector. Recently the Sobolev differentiability of the solution to an SDE was proved in [3, 4, 6, 7] under rather weak assumptions on the drift. The methods of these works differ from ours. In particular, the Malliavin calculus is used in [6, 7]. The condition we impose on the drift is more restrictive than that of [3, 7], but it allows us to obtain an explicit representation for the derivative in terms of intrinsic parameters of the initial equation. This representation is a natural generalization of the expression for the derivative in the smooth case.

In the one-dimensional case [1, 2] the derivative was represented via the local time of the process. It is well known that the solution does not have a local time at a point in multidimensional situation. We use continuous additive functionals for the representation of the derivative. This method can be considered as a generalization of the local time approach to the multidimensional case.

We show that the derivative is a solution of an integral equation. Our method is likely to can be used in the case of non-constant diffusion.

REFERENCES

- [1] O. V. Aryasova and A. Yu. Pilipenko. On properties of a flow generated by an SDE with discontinuous drift. *Electron. J. Probab.*, 17:no. 106, 1–20, 2012.
- [2] S. Attanasio. Stochastic flows of diffeomorphisms for one-dimensional SDE with discontinuous drift. *Electron. Commun. Probab.*, 15:no. 20, 213–226, 2010.
- [3] E. Fedrizzi and F. Flandoli. Hölder flow and differentiability for SDEs with nonregular drift. *Stochastic Analysis and Applications*, 31(4):708–736, 2013.
- [4] E. Fedrizzi and F. Flandoli. Noise prevents singularities in linear transport equations. *Journal of Functional Analysis*, 264(6):1329 – 1354, 2013.
- [5] F. Flandoli, M. Gubinelli, and E. Priola. Flow of diffeomorphisms for SDEs with unbounded Hölder continuous drift. *Bulletin des Sciences Mathématiques*, 134(4):405 – 422, 2010.
- [6] T. Meyer-Brandis and F. Proske. Construction of strong solutions of SDE’s via Malliavin calculus. *Journal of Functional Analysis*, 258(11):3922 – 3953, 2010.
- [7] S.E.A. Mohammed, T. Nilssen, and F. Proske. Sobolev differentiable stochastic flows of SDE’s with measurable drift and applications. 2012. arXiv/1204.3867.
- [8] A. Y. Veretennikov. On strong solutions and explicit formulas for solutions of stochastic integral equations. *Math. USSR Sborn*, 39(3):387–403, 1981.

INSTITUTE OF GEOPHYSICS, NATIONAL ACADEMY OF SCIENCES OF UKRAINE, PALLADIN PR.,
32, KIEV-142, UKRAINE

E-mail address: oaryasova@gmail.com